

ordered
ref. 4/18

> s ovarian cancer

68566 OVARIAN
314596 CANCER
L1 9941 OVARIAN CANCER
(OVARIAN(W) CANCER)

=> s pi kinase

30449 PI
129484 KINASE
L2 136 PI KINASE
(PI(W) KINASE)

=> s l1 and l2

L3 0 L1 AND L2

=> s pik3ca

L4 6 PIK3CA

=> s l1 and l4

L5 3 L1 AND L4

=> d l5 1-3 bib,ab

L5 ANSWER 1 OF 3 MEDLINE
AN 2000090217 MEDLINE
DN 20090217
TI Growth suppression of human **ovarian cancer** cells by
adenovirus-mediated transfer of the PTEN gene.
AU Minaguchi T; Mori T; Kanamori Y; Matsushima M; Yoshikawa H; Taketani Y;
Nakamura Y
CS Laboratory of Molecular Medicine, Human Genome Center, The Institute of
Medical Science, The University of Tokyo, Japan.
SO CANCER RESEARCH, (1999 Dec 15) 59 (24) 6063-7.
Journal code: CNF. ISSN: 0008-5472.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 200003
EW 20000305
AB A tumor suppressor gene on chromosome 10q23, PTEN, encodes a
phosphatidylinositol phosphatase that antagonizes activation of the
phosphatidylinositol 3'-kinase-mediated pathway involved in cell growth.
A gene encoding the catalytic subunit of phosphatidylinositol 3'-kinase (
PIK3CA) is frequently activated in ovarian cancers; therefore,
overexpression of the PTEN product through gene transfer might be an
effective strategy for treating ovarian cancers. To test the potential
for this type of gene therapy, we constructed a recombinant adenovirus
encoding wild-type PTEN and examined its effects on nine cell lines
derived from human ovarian carcinomas. Transduction of the PTEN gene
significantly inhibited growth of six of these cell lines compared with

infection with virus alone, and the degree of inhibition correlated with the efficiency of gene transfer as determined by β -galactosidase assay.

Results of flow cytometry suggested that the observed effects were mediated by two mechanisms, apoptosis and/or arrest in the G1 phase of the

cell cycle, and that high adenoviral transduction efficiency of cells was associated with induction of apoptosis. We also found that the level of transcription of Integrin α (v) in **ovarian cancer** cells correlated with the efficiency of transduction ($P = 0.014$) and with the degree of growth inhibition after PTEN gene transfer ($P = 0.009$). These findings carry significant implications for adenovirus vector-based PTEN gene therapies for ovarian cancers.

L5 ANSWER 2 OF 3 MEDLINE

AN 2000025001 MEDLINE

DN 20025001

TI **PIK3CA**: determining its role in cellular proliferation and **ovarian cancer**.

AU Andrew S

CS Department of Medical Genetics, University of Alberta, Edmonton, Canada..
seandrew@pop.srv.ualberta.ca

SO CLINICAL GENETICS, (1999 Sep) 56 (3) 190-1.
Journal code: DDT. ISSN: 0009-9163.

CY Denmark

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200001

EW 20000104

L5 ANSWER 3 OF 3 MEDLINE

AN 1999113837 MEDLINE

DN 99113837

TI **PIK3CA** is implicated as an oncogene in **ovarian cancer** [see comments].

CM Comment in: Nat Genet 1999 Jan;21(1):64-5

AU Shayesteh L; Lu Y; Kuo W L; Baldocchi R; Godfrey T; Collins C; Pinkel D; Powell B; Mills G B; Gray J W

CS UCSF Cancer Center, University of California, San Francisco 941430-0808, USA.

NC CA09215 (NCI)

P01-CA64602 (NCI)

SO NATURE GENETICS, (1999 Jan) 21 (1) 99-102.

Journal code: BRO. ISSN: 1061-4036.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199904

EW 19990402

AB **Ovarian cancer** is the leading cause of death from gynecological malignancy and the fourth leading cause of cancer death among American women, yet little is known about its molecular aetiology. Studies using comparative genomic hybridization (CGH) have revealed several regions of recurrent, abnormal, DNA sequence copy number that may encode genes involved in the genesis or progression of the disease. One region at 3q26 found to be increased in copy number in approximately 40% of ovarian and others cancers contains **PIK3CA**, which encodes the p110 α catalytic subunit of phosphatidylinositol 3-kinase (PI3-kinase).

The association between **PIK3CA** copy number and PI3-kinase activity makes **PIK3CA** a candidate oncogene because a broad range of cancer-related functions have been associated with PI3-kinase mediated signalling. These include proliferation, glucose transport and catabolism,

cell adhesion, apoptosis, RAS signalling and oncogenic transformation. In addition, downstream effectors of PI3-kinase, AKT1 and AKT2, have been found to be amplified or activated in human tumours, including ovarian cancer. We show here that **PIK3CA** is frequently increased in copy number in ovarian cancers, that the increased copy number is associated with increased **PIK3CA** transcription, p110alpha protein expression and PI3-kinase activity and that treatment with the PI3-kinase inhibitor LY294002 decreases proliferation and increases apoptosis. Our observations suggest **PIK3CA** is an oncogene that has an important role in ovarian cancer.

N 2000090217 MEDLINE
 DN 20090217
 TI Growth suppression of human **ovarian cancer** cells by
 adenovirus-mediated transfer of the PTEN gene.
 AU Minaguchi T; Mori T; Kanamori Y; Matsushima M; Yoshikawa H; Taketani Y;
 Nakamura Y
 CS Laboratory of Molecular Medicine, Human Genome Center, The Institute of
 Medical Science, The University of Tokyo, Japan.
 SO CANCER RESEARCH, (1999 Dec 15) 59 (24) 6063-7.
 Journal code: CNF. ISSN: 0008-5472.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals; Cancer Journals
 EM 200003
 EW 20000305
 AB A tumor suppressor gene on chromosome 10q23, PTEN, encodes a
 phosphatidylinositol phosphatase that antagonizes activation of the
phosphatidylinositol 3'-kinase-mediated pathway involved
 in cell growth. A gene encoding the catalytic subunit of
phosphatidylinositol 3'-kinase (PIK3CA) is frequently
 activated in ovarian cancers; therefore, overexpression of the PTEN
 product through gene transfer might be an effective strategy for treating
 ovarian cancers. To test the potential for this type of gene therapy, we
 constructed a recombinant adenovirus encoding wild-type PTEN and examined
 its effects on nine cell lines derived from human ovarian carcinomas.
 Transduction of the PTEN gene significantly inhibited growth of six of
 these cell lines compared with infection with virus alone, and the degree
 of inhibition correlated with the efficiency of gene transfer as
 determined by beta-galactosidase assay. Results of flow cytometry
 suggested that the observed effects were mediated by two mechanisms,
 apoptosis and/or arrest in the G1 phase of the cell cycle, and that high
 adenoviral transduction efficiency of cells was associated with induction
 of apoptosis. We also found that the level of transcription of Integrin
 alpha(v) in **ovarian cancer** cells correlated with the
 efficiency of transduction ($P = 0.014$) and with the degree of growth
 inhibition after PTEN gene transfer ($P = 0.009$). These findings carry
 significant implications for adenovirus vector-based PTEN gene therapies
 for ovarian cancers.

L7 ANSWER 2 OF 7 MEDLINE
 AN 2000025001 MEDLINE
 DN 20025001
 TI PIK3CA: determining its role in cellular proliferation and **ovarian**
cancer.
 AU Andrew S
 CS Department of Medical Genetics, University of Alberta, Edmonton, Canada..
 seandrew@pop.srv.ualberta.ca
 SO CLINICAL GENETICS, (1999 Sep) 56 (3) 190-1.
 Journal code: DDT. ISSN: 0009-9163.
 CY Denmark
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200001
 EW 20000104

L7 ANSWER 3 OF 7 MEDLINE
 AN 1999184222 MEDLINE

DN 99184222
TI **Ovarian cancer** investigators aim at cell signaling pathways [news].
AU Friedrich M J
SO JAMA, (1999 Mar 17) 281 (11) 973-5.
Journal code: KFR. ISSN: 0098-7484.
CY United States
DT News Announcement
LA English
FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals
EM 199905
EW 19990504

L7 ANSWER 4 OF 7 MEDLINE
AN 1999113837 MEDLINE
DN 99113837
TI PIK3CA is implicated as an oncogene in **ovarian cancer** [see comments].
CM Comment in: Nat Genet 1999 Jan;21(1):64-5
AU Shayesteh L; Lu Y; Kuo W L; Baldocchi R; Godfrey T; Collins C; Pinkel D; Powell B; Mills G B; Gray J W
CS UCSF Cancer Center, University of California, San Francisco 941430-0808, USA.
NC CA09215 (NCI)
P01-CA64602 (NCI)
SO NATURE GENETICS, (1999 Jan) 21 (1) 99-102.
Journal code: BRO. ISSN: 1061-4036.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199904
EW 19990402
AB **Ovarian cancer** is the leading cause of death from gynecological malignancy and the fourth leading cause of cancer death among American women, yet little is known about its molecular aetiology. Studies using comparative genomic hybridization (CGH) have revealed several regions of recurrent, abnormal, DNA sequence copy number that may encode genes involved in the genesis or progression of the disease. One region at 3q26 found to be increased in copy number in approximately 40% of ovarian and others cancers contains PIK3CA, which encodes the p110alpha catalytic subunit of **phosphatidylinositol 3-kinase** (PI3-kinase). The association between PIK3CA copy number and PI3-kinase activity makes PIK3CA a candidate oncogene because a broad range of cancer-related functions have been associated with PI3-kinase mediated signalling. These include proliferation, glucose transport and catabolism, cell adhesion, apoptosis, RAS signalling and oncogenic transformation. In addition, downstream effectors of PI3-kinase, AKT1 and AKT2, have been found to be amplified or activated in human tumours, including **ovarian cancer**. We show here that PIK3CA is frequently increased in copy number in ovarian cancers, that the increased copy number is associated with increased PIK3CA transcription, p110alpha protein expression and PI3-kinase activity and that treatment with the PI3-kinase inhibitor LY294002 decreases proliferation and increases apoptosis. Our observations suggest PIK3CA is an oncogene that has an important role in **ovarian cancer**.

L7 ANSWER 5 OF 7 MEDLINE
AN 97159701 MEDLINE
DN 97159701
TI HER-2/neu signal transduction in human breast and **ovarian cancer**.
AU Reese D M; Slamon D J

CS Division of Hematology/Oncology and Jonsson Comprehensive Cancer Center,
 UCLA School of Medicine, Los Angeles, California 90095, USA.
 NC 1K12CA01714 (NCI)
 R01 CA36827 (NCI)
 SO STEM CELLS, (1997) 15 (1) 1-8. Ref: 85
 Journal code: BN2. ISSN: 1066-5099.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 General Review; (REVIEW)
 (REVIEW, TUTORIAL)
 LA English
 FS Priority Journals
 EM 199707
 AB The HER-2/neu proto-oncogene encodes a 185 kDa transmembrane receptor
 tyrosine kinase with significant sequence homology to other members of
 the class I receptor tyrosine kinase family. The HER-2/neu gene is amplified
 and/or overexpressed in 25%-30% of human breast and ovarian cancers, and
 overexpression of the receptor is associated with poor prognosis.

Tyrosine phosphorylation and activation of the HER-2 receptor lead to activation
 of specific signal transduction pathways in breast and **ovarian**
cancer cells, including the ras/MAP **kinase** cascade,
phosphatidylinositol 3-kinase, and phospholipase
 C-gamma. HER-2/neu signal transduction pathways ultimately converge on
 the cell nucleus, where the expression of diverse genes is induced after
 activation of the receptor. A more complete understanding of HER-2/neu
 signal transduction pathways may allow the development of specific
 therapeutics for the treatment of those human breast and ovarian cancers
 containing this alteration.

L7 ANSWER 6 OF 7 MEDLINE
 AN 96150936 MEDLINE
 DN 96150936
 TI SH2 and SH3 domains: potential targets for anti-cancer drug design.
 AU Smithgall T E
 CS Eppley Institute for Research in Cancer, University of Nebraska Medical
 Center, Omaha 68198-6805, USA.
 SO JOURNAL OF PHARMACOLOGICAL AND TOXICOLOGICAL METHODS, (1995 Nov) 34 (3)
 125-32. Ref: 52
 Journal code: A9W. ISSN: 1056-8719.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 General Review; (REVIEW)
 (REVIEW, TUTORIAL)
 LA English
 FS Priority Journals
 EM 199605
 AB Protein-tyrosine kinases interact with a diverse group of signaling
 molecules that share common structural elements known as Src homology 2
 and 3 (SH2 and SH3) domains. SH2 domains bind with high affinity to
 peptide sequences within target proteins that contain phosphorylated
 tyrosine residues, but have no affinity for the unphosphorylated
 sequence.
 This property allows activated tyrosine kinases to initiate signal
 transduction by recruiting downstream effectors with SH2 domains. SH3
 domains also mediate protein-protein interaction. Target sequences for
 SH3 domains are rich in proline and hydrophobic amino acids, but do not
 require phosphorylation. SH2- and SH3-mediated protein-protein
 interactions are required for the transmission of proliferative signals
 initiated by tyrosine kinases (e.g., Ras activation or stimulation of
phosphatidylinositol-3' kinase activity). Peptidomimetic

ligands based on the sequence of target proteins for SH2 and SH3 domains may represent new lead compounds for the therapy of proliferative diseases

that are dependent upon constitutively activated tyrosine kinases (e.g., BCR/ABL in chronic myelogenous and acute lymphocytic leukemias or HER-2/Neu in breast and **ovarian cancer**).

L7 ANSWER 7 OF 7 MEDLINE

AN 95309792 MEDLINE

DN 95309792

TI Evidence for tight coupling of gonadotropin-releasing hormone receptors to

phosphatidylinositol kinase in plasma membrane from ovarian carcinomas.

AU Takagi H; Imai A; Furui T; Horibe S; Fuseya T; Tamaya T

CS Department of Obstetrics and Gynecology, Gifu University School of Medicine, Japan.

SO GYNECOLOGIC ONCOLOGY, (1995 Jul) 58 (1) 110-5.

Journal code: FXC. ISSN: 0090-8258.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals; Cancer Journals

EM 199509

AB Gonadotropin-releasing hormone (Gn-RH) analogs inhibit **ovarian cancer** cell proliferation in vivo and in vitro. To examine whether Gn-RH receptor (Gn-RHR) mediates direct antiproliferative effects, we attempted to determine inhibitory regulation by Gn-RH of **phosphatidylinositol** (PtdIns) **kinase** activity, known to stimulate mitogenic response, in plasma membranes isolated from ovarian carcinoma samples. Ovarian carcinomas surgically removed and cloned cell line SK-OV3 had been screened for Gn-RHR expression prior to plasma membrane isolation. PtdIns kinase activity was measured as

phosphorylation

of exogenous substrate PtdIns by the purified plasma membranes.

Incubation

of the plasma membranes isolated from Gn-RHR-positive specimens with [γ -³²P]ATP and PtdIns caused [³²P]phosphate incorporation into PtdIns phosphate (PtdInsP) in a time-dependent manner. Concomitant exposure of the membrane preparations to Gn-RH analog buserelin (1 microM) led to a 70% inhibition of the PtdInsP production, when compared to control. After 10 or 15 min of an initial incubation, the addition of analog resulted in similar suppression of PtdIns phosphorylation. This inhibition was dependent on the buserelin dose, and a half-maximal effect occurred at a concentration 0.1 to 1 nM of buserelin. Degradation of the produced PtdInsP in the plasma membranes was not affected by the Gn-RH analog. Similar inhibition of PtdIns kinase activities was observed in membranes prepared from cells that had been pretreated with buserelin (1 microM)

for

48 hr prior to assay. These findings demonstrate that PtdIns kinase activity is suppressed by Gn-RH analog in plasma membrane isolated from GnRHR-expressing ovarian carcinomas, suggesting a tight coupling of

Gn-RHR

to PtdIns. The inhibition of membrane-associated PtdIns kinase by Gn-RHR occupancy may mediate the antimitogenic action of the hormone on human ovarian carcinomas.

8 ANSWER 3 OF 4 MEDLINE
AN 1999422308 MEDLINE
DN 99422308
TI Expression analysis of genes at 3q26-q27 involved in frequent
amplification in squamous cell lung carcinoma.
AU Racz A; Brass N; Heckel D; Pahl S; Remberger K; Meese E
CS Department of Human Genetics, Medical School, University of Saarland,
Homburg/Saar, Germany.
SO EUROPEAN JOURNAL OF CANCER, (1999 Apr) 35 (4) 641-6.
Journal code: ARV. ISSN: 0959-8049.
CY ENGLAND: United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199912
EW 19991202
AB Gene amplifications are known to occur frequently in lung **cancer**
. Recently, we identified gene amplifications at 3q26 in squamous cell
lung carcinoma (SCC) using reverse chromosome painting. Here, our aim was
to analyse the expression of genes which map within the amplified
chromosomal region. The genes which were selected for their known
function
and their potential involvement in tumour development included the genes
for ribosomal protein L22 (RPL22), butyrylcholinesterase (BCHE), glucose
transporter 2 (SLC2A2), transferrin receptor (TFRC), thrombopoietin
(THPO)
and the phosphatidylinositol-3 kinase catalytic alpha polypeptide (**PIK3CA**). While five genes were expressed in the majority of the 17
samples of SCC, the gene for the glucose transporter 2 (SLC2A2) was
expressed in only three cases, excluding SLC2A2 as the target gene of the
amplification unit. For a subset of tumours, we determined the
amplification status of the six genes. The TFRC, **PIK3CA**, BCHE,
THPO and SLC2A2 genes were amplified in several cases, whereas the RPL22
gene was amplified in only one case. The combined amplification and
expression data of this and our previous studies indicate that the
amplified region at 3q26 contains several genes that are transcribed in
SCC, providing the possibility that several amplified and functionally
important genes at 3q26 may be involved in the pathogenesis of SCC.